

FDA Safe Food Handling

Safe steps in food handling, cooking, and storage are essential to prevent foodborne illness. You can't see, smell, or taste harmful bacteria that may cause illness. In every step of food preparation, follow the four Fight BAC!™ guidelines to keep food safe:

- Clean — Wash hands and surfaces often.
- Separate — Don't cross-contaminate.
- Cook — Cook to proper temperatures.
- Chill — Refrigerate promptly.

Shopping

- Purchase refrigerated or frozen items after selecting your non-perishables.
- Never choose meat or poultry in packaging that is torn or leaking.
- Do not buy food past "Sell-By," "Use-By," or other expiration dates.

Storage

- Always refrigerate perishable food within 2 hours (1 hour when the temperature is above 90 °F).
- Check the temperature of your refrigerator and freezer with an appliance thermometer. The refrigerator should be at 40 °F or below and the freezer at 0 °F or below.
- Cook or freeze fresh poultry, fish, ground meats, and variety meats within 2 days; other beef, veal, lamb, or pork, within 3 to 5 days.
- Perishable food such as meat and poultry should be wrapped securely to maintain quality and to prevent meat juices from getting onto other food.
- To maintain quality when freezing meat and poultry in its original package, wrap the package again with foil or plastic wrap that is recommended for the freezer.
- In general, high-acid canned food such as tomatoes, grapefruit, and pineapple can be stored on the shelf for 12 to

18 months. Low-acid canned food such as meat, poultry, fish, and most vegetables will keep 2 to 5 years — if the can remains in good condition and has been stored in a cool, clean, and dry place. Discard cans that are dented, leaking, bulging, or rusted.

Preparation

- Always wash hands with warm water and soap for 20 seconds before and after handling food.
- Don't cross-contaminate. Keep raw meat, poultry, fish, and their juices away from other food. After cutting raw meats, wash cutting board, utensils, and countertops with hot, soapy water.
- Cutting boards, utensils, and countertops can be sanitized by using a solution of 1 tablespoon of unscented, liquid chlorine bleach in 1 gallon of water.
- Marinate meat and poultry in a covered dish in the refrigerator.

Thawing

- **Refrigerator:** The refrigerator allows slow, safe thawing. Make sure thawing meat and poultry juices do not drip onto other food.
- **Cold Water:** For faster thawing, place food in a leak-proof plastic bag. Submerge in cold tap water. Change the water every 30 minutes. Cook immediately after thawing.
- **Microwave:** Cook meat and poultry immediately after microwave thawing.

Cooking

- Cook all raw beef, pork, lamb and veal steaks, chops, and roasts to a minimum internal temperature of 145 °F as measured with a food thermometer before removing meat from the heat source. For safety and quality, allow meat to rest for at least three minutes before carving or consuming. For reasons of personal preference, consumers may choose to cook meat to higher temperatures.

Ground meats: Cook all raw ground beef, pork, lamb, and

veal to an internal temperature of 160 °F as measured with a food thermometer.

Poultry: Cook all poultry to an internal temperature of 165 °F as measured with a food thermometer.

Serving

- Hot food should be held at 140 °F or warmer.
- Cold food should be held at 40 °F or colder.
- When serving food at a buffet, keep food hot with chafing dishes, slow cookers, and warming trays. Keep food cold by nesting dishes in bowls of ice or use small serving trays and replace them often.
- Perishable food should not be left out more than 2 hours at room temperature (1 hour when the temperature is above 90 °F).

Leftovers

- Discard any food left out at room temperature for more than 2 hours (1 hour if the temperature was above 90 °F).
- Place food into shallow containers and immediately put in the refrigerator or freezer for rapid cooling.
- Use cooked leftovers within 4 days.
- Reheat leftovers to 165 °F.

Refreezing

Meat and poultry defrosted in the refrigerator may be refrozen before or after cooking. If thawed by other methods, cook before refreezing.